



Missouri Pharmacy Program – Preferred Drug List



Tramadol-like Agents

Effective 06/21/2012

Revised 04/02/2015

Preferred Agents

- Tramadol
- Tramadol-Apap

Non-Preferred Agents

- Ultram®
- Ultracet®
- Ultram ER®
- Tramadol ER
- Ryzolt ER®/ODT
- Nucynta®
- NucyntaER®
- Conzip®
- Rybix® ODT

<u>Approval Criteria</u>	<u>Denial Criteria</u>
Failure to achieve desired therapeutic outcomes with trial on 2 or more preferred agents <ul style="list-style-type: none">• Documented trial period for preferred agents• Documented ADE/ADR to preferred agents	(see below)
Documented compliance on current therapy regimen	Drug Prior Authorization Hotline: (800) 392-8030

Denial Criteria

Ultram:

- 15 years of age or less²
- Daily dose exceeds maximum recommended daily dose:
 - Patients \leq 75 years of age – 400 mg
 - Patients $>$ 75 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

Ultracet:

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
 - Patients \geq 16 years of age – 300 mg tramadol and 2600mg acetaminophen

- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

Ultram ER/Ryzolt:

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
 - Patients \geq 16 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days
- History of opioid abuse or dependence – therapy will be subject to Clinical Consultant review
- Claim for Ultram ER/Ryzolt therapy without previous stabilization on immediate-release tramadol product.

Nucynta:

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
 - Patients \geq 18 years of age – 700 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

Nucynta ER:

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
 - Patients \geq 18 years of age – 500 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

Rybix ODT

- 16 years of age or less
- Daily dose exceeds maximum recommended daily dose:
 - Patients \geq 17 years of age – 400mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days